

Infection Protocol for Recombinant Lentivirus

--- Storage Instructions and Infection Protocol



9601 Medical Center Drive
A&R Building, Suite 341
Rockville, MD 20850
FAX. 301-560-4919
TEL. 301-330-5966
Toll Free. 1-(866)-918-6812
Email: info@signagen.com
Web: www.signagen.com

This product is for laboratory research ONLY and not for diagnostic use

Introduction:

Lentivirus is a genus of viruses of the retroviridae family and has the unique ability among retroviruses of being able to infect both dividing non-dividing cells efficiently. Lentivirus is one of the most efficient vector to transduce mammalian cells for both in vitro and in vivo.

Important Guidelines for Transduction:

- The optimal MOI (multiplicity of infection) is target cell specific. To determine the optimal MOI, it is advisable to perform several test transductions with reporter viral particles (e.g. [LV-CMV-GFP](#) or [LV-CMV-mCherry](#)) in different MOIs at 0.5, 2, 8, 20, 40 etc. The optimal MOI is then chosen to yield the highest percentage of successfully infected cells without invisible toxicity.
- Infection with the presence of hexadimethrine bromide (polybrene) at final 3-8 µg/mL in the cell culture medium enhances transduction of most cell types by 2-10 fold.
- The Infection Protocol for Suspension Cell is recommended for transduction of suspension cells and includes an additional spin step which concentrates the virus at the target cell surface to increase transduction efficiency.
- Transduction efficiency should be checked ~72 hrs following infection if no antibiotic selection is used.
- For cells that are more resistant to infection, it is desirable to select only the clones that stably express the transgene for downstream biological assays.

In-vitro Infection Protocol for Adherent Cell:

Step 1. Prepare Virus-containing Media:

- Thaw the lentiviral stock at RT.
- Add desired amount of virus to growth media to achieve the desired optimal MOI. Lentiviral transduction unit (TU) to be used = MOI x number of cells to be infected. For example, if you intend to infect 50 thousands cells at MOI of 10, you need $10 \times 50,000 = 5.0 \times 10^5$ TU for the infection. If the original lentivirus stock is 10^9 TU/mL, then you will need 0.5 µL of the original stock for the dilution.

Step 2. Infecting Cells with Lentivirus:

- Remove the original cell culture media.
- Add the above lentivirus-containing media prepared from **Step 1** to cell culture. Below is a general guideline for the amount of media used:

24-well plate:	0.2-0.3 mL
12-well plate:	0.5-0.8 mL
6-well plate:	1-1.5 mL/well
60 mm-plate:	3-4 mL/plate
10 cm-plate:	8-12 mL/plate

- Optional: Add polybrene at final 3-8 µg/mL to cell culture medium enhances transduction of most cell types by 2-10 fold.
- Incubate cells with the virus-containing media for overnight followed by removing virus-containing media and replacing with fresh, desired media next day after infection.
- Check transduction efficiency 3-5 days after infection.
- To create a stable cell line, add appropriate antibiotics 24-48 hrs post transduction.

In-vitro Infection Protocol for Suspension Cell:

The following protocol has been given as a general starting point for 6-well plate. The transduction conditions shall be scaled up or down per the surface of tissue culture dishes.

- Resuspend the target cell to final 8 mL at density of $\sim 10^6$ cells/mL. Aliquot 2 mL into each of four 15 mL sterile conical tubes.
- Add appropriate volume of virus at optimal MOI to infect cells with the presence of polybrene at final 8 µg/mL. A reporter lentivirus like [LV-CMV-GFP](#) or [LV-CMV-mCherry](#) shall be included to monitor the transduction efficiency.
- Gently mix and incubate cells for 10 min at RT.
- Centrifuge cells for 30 min at 800xg at 32 °C.
- Remove virus containing medium and resuspend cell pellet with 2 mL of fresh complete culture media followed by transfer of 2 mL cell suspension to a well of 6-well plate.
- Incubate cells for 24-72 hrs.
- Check transduction efficiency 3-5 days after infection.
- Split cells 1:3 or 1:5 and continue incubation for 48 hrs in complete media.
- The infected cells can then be selected for stable expression using appropriate antibiotic selection.
- Biological assays can be then carried out.

Storage: Lentivirus stock is supplied in sterile PBS at indicated titer. The lentivirus can be stored at -80 °C 24 months without a significant loss in titer. If desired, aliquot viral stock upon arrival and store those aliquots at -80 °C freezer immediately. **DO NOT FREEZE AND THAW REPEATEDLY.**

Infection Protocol of Recombinant Lentivirus)

- Storage Instructions and Infection Protocol



9601 Medical Center Drive

A&R Building, Suite 341

Rockville, MD 20850

FAX. 301-560-4919

TEL. 301-330-5966

Toll Free. 1-(866)-918-6812

Email: info@signagen.com

Web: www.signagen.com

This product is for laboratory research ONLY and not for diagnostic use

***In-vivo* Infection Protocol:**

The following protocol is given for stereotaxic gene delivery of lentivirus to the rodent brain.

Materials:

- Super purified lentivirus in PBS, in vivo grade.
- Ethanol 70%.
- Mice or rats.
- Anesthetics and analgesics (e.g., ketamine etc).
- Sterile PBS
- Bone wax
- Triple antibiotic ointment

Procedures:

1. Anesthetize and fix animal in stereotaxic apparatus.
2. Make incision and locate bregma.
3. Surgery - Preparation of craniotomy by referring to the standard stereotaxic coordinates of mouse or rats. The craniotomy is drilled using a hand-held drill. The following table is an examples of stereotaxic coordinates.

Targeted region adult mouse brain00	Rostral (+) caudal (-)(mm)	Lateral (mm)	Ventral (mm)
Subthalamic nucleus	- 1.9	1.6	4.4
Dorsal hippocampus, CA1	- 2.1	2.0	1.4
Basolateral amygdala	- 1.5	2.75	4.75
Lateral ventricle	+ 0.5	0.75	2.5
Nucleus accumbens, core	+ 1.1	1.2	4.5

4. Injection of lentivirus for a single injection. Place the injection micropipette into the holder of the stereotaxic arm. Fill the micropipette with 3 μ L lentivirus. Then slowly lower the micropipette to the desired z coordinate of the injection site. Apply pressure steadily via a pump or syringe to inject 100~500 nL lentivirus at $>1E+9$ TU/mL within 60~90 seconds.
5. After recovery of animal, feed the animal as usual until the transgene expression is checked one week after lentivirus injection.

Note: If lentivirus is administrated to mouse via different routes, please refer the lentivirus injection dose to the following table.

Lentivirus Administration Route	Dose (TU)
Intracerebral	$5 \times 10^5 \sim 5 \times 10^6$
Intraventricular	$10^6 \sim 5 \times 10^6$
Intravenous	$\sim 5 \times 10^8$
Intranasal	$10^7 \sim 3 \times 10^7$
Intravitreal	$10^6 \sim 4 \times 10^6$